

GEOMETRIC CONVEXITY OF THE GENERALIZED SINE AND THE GENERALIZED HYPERBOLIC SINE

WEI-DONG JIANG AND FENG QI

ABSTRACT. In the paper, the authors prove that the generalized sine function $\sin_{p,q}(x)$ and the generalized hyperbolic sine function $\sinh_{p,q}(x)$ are geometrically concave and geometrically convex, respectively. Consequently, the authors verify a conjecture posed in the paper “B. A. Bhayo and M. Vuorinen, *On generalized trigonometric functions with two parameters*, J. Approx. Theory **164** (2012), no. 10, 1415–1426; Available online at <http://dx.doi.org/10.1016/j.jat.2012.06.003>”.

1. INTRODUCTION

It is well known from calculus that

$$\arcsin x = \int_0^x \frac{1}{(1-t^2)^{1/2}} dt$$

for $0 \leq x \leq 1$ and

$$\frac{\pi}{2} = \arcsin 1 = \int_0^1 \frac{1}{(1-t^2)^{1/2}} dt.$$

We can define the sine function on $[0, \frac{\pi}{2}]$ as the inverse of the arcsine function and extend it to $(-\infty, \infty)$.

Let $1 < p < \infty$. the arcsine can be generalized as

$$\arcsin_p x = \int_0^x \frac{1}{(1-t^p)^{1/p}} dt, \quad 0 \leq x \leq 1$$

and

$$\frac{\pi_p}{2} = \arcsin_p 1 = \int_0^1 \frac{1}{(1-t^p)^{1/p}} dt.$$

The inverse of the function \arcsin_p on $[0, \frac{\pi_p}{2}]$ is called the generalized sine function and denoted by \sin_p . By standard extending procedures as the sine function done, we may obtain a differentiable function on the whole real line $(-\infty, \infty)$, which coincides with the sine when $p = 2$. It is easy to see that the function \sin_p is strictly increasing and concave on $[0, \frac{\pi_p}{2}]$. Similarly, we can define the generalized cosine, tangent, hyperbolic functions and their inverses.

For $p, q > 1$, let

$$F_{p,q}(x) = \int_0^x (1-t^q)^{-1/p} dt, \quad x \in [0, 1]$$

2010 *Mathematics Subject Classification*. Primary 33B10; Secondary 26A51, 26D05, 33C20.

Key words and phrases. generalized sine, generalized hyperbolic sine, geometric convexity, inequality, conjecture.

This work was supported in part by the Project of Shandong Province Higher Educational Science and Technology Program under grant No. J11LA57.

and

$$\frac{\pi_{p,q}}{2} = \int_0^1 (1-t^q)^{-1/p} dt.$$

Then $F_{p,q} : [0, 1] \rightarrow [0, \frac{\pi_{p,q}}{2}]$ is an increasing homeomorphism. We denote this function $F_{p,q}$ by $\arcsin_{p,q}$. Thus, its inverse

$$\sin_{p,q} = F_{p,q}^{-1}$$

is defined on the interval $[0, \frac{\pi_{p,q}}{2}]$. By a similar extension to the sine function, we can find a differentiable function $\sin_{p,q}$ defined on \mathbb{R} . We call $\sin_{p,q}$ the generalized (p, q) -sine function.

We can also define

$$\arccos_{p,q} x = \arcsin_{p,q} [(1-x^p)^{1/q}],$$

see [2, 4], and the inverse of the generalized (p, q) -hyperbolic sine function

$$\operatorname{arcsinh}_{p,q}(x) = \int_0^x (1+t^q)^{-1/p} dt, \quad x \in (0, \infty).$$

Their inverse functions are

$$\sin_{p,q} : \left(0, \frac{\pi_{p,q}}{2}\right) \rightarrow (0, 1), \quad \cos_{p,q} : \left(0, \frac{\pi_{p,q}}{2}\right) \rightarrow (0, 1),$$

and

$$\sinh_{p,q} : (0, m_{p,q}^*) \rightarrow (0, \infty),$$

where

$$m_{p,q}^* = \int_0^\infty (1+t^q)^{-1/p} dt.$$

When $p = q$, the (p, q) -functions $\sin_{p,q}$, $\cos_{p,q}$, $\sinh_{p,q}$, $\arcsin_{p,q}$, $\arccos_{p,q}$, and $\operatorname{arcsinh}_{p,q}$ reduce to p -functions \sin_p , \cos_p , \sinh_p , \arcsin_p , \arccos_p , and $\operatorname{arcsinh}_p$ respectively. See [3, 5, 7]. In particular, when $p = q = 2$, the (p, q) -functions become our familiar trigonometric and hyperbolic functions.

Recently, the generalized trigonometric and hyperbolic functions, including (p, q) -functions and p -functions, have been studied by many mathematicians from different points of view. See [1, 6, 8, 11]. In [4], the authors gave basic properties of the generalized (p, q) -trigonometric functions. In [5], the authors generalized some classical inequalities for trigonometric and hyperbolic functions, such as Mitri-nović-Adamović inequality, Lazarević's inequality, Huygens-type inequalities, and Wilker-type inequalities.

In [2], the authors found that the functions $\arcsin_{p,q}$ and $\operatorname{arcsinh}_{p,q}$ can be expressed in terms of Gaussian hypergeometric functions. Applying the vast available information about hypergeometric functions, some remarkable properties and inequalities for generalized trigonometric and hyperbolic functions are obtained. Moreover, they raised the following conjecture.

Conjecture 1.1 ([2, p. 1421, Conjecture 2.11]). *If $p, q \in (1, \infty)$ and $r, s \in (0, 1)$, then*

$$\sin_{p,q} \sqrt{rs} \geq \sqrt{\sin_{p,q} r \sin_{p,q} s} \tag{1.1}$$

and

$$\sinh_{p,q} \sqrt{rs} \leq \sqrt{\sinh_{p,q} r \sinh_{p,q} s}. \tag{1.2}$$

The main purpose of this paper is to discover the geometric convexity of $\sin_{p,q}(x)$ and $\sinh_{p,q}(x)$ and to give an affirmative answer to the above stated [2, p. 1421, Conjecture 2.11].

Our main results may be formulated in the following theorem.

Theorem 1.1. *Let $p, q \in (1, \infty)$ and $r, s, x \in (0, 1)$. Then*

- (1) *the function $\sin_{p,q}(x)$ is geometrically concave function;*
- (2) *the function $\sinh_{p,q}(x)$ is geometrically convex function.*

Corollary 1.1. *Let $p, q \in (1, \infty)$ and $r, s, x \in (0, 1)$. Then the inequalities (1.1) and (1.2) and the inequalities*

$$\sin_p \sqrt{rs} \geq \sqrt{\sin_p r \sin_p s} \quad (1.3)$$

and

$$\sinh_p \sqrt{rs} \leq \sqrt{\sinh_p r \sinh_p s} \quad (1.4)$$

are all valid.

2. A DEFINITION AND LEMMAS

For proving our main results, we need the following definition and lemmas.

Definition 2.1. Let $I \subseteq (0, +\infty)$ be an interval and $f : I \rightarrow (0, \infty)$ be a continuous function. This function f is said to be geometrically convex on I if

$$f(x^\lambda y^{1-\lambda}) \leq f^\lambda(x) f^{1-\lambda}(y). \quad (2.1)$$

is valid for all $x, y \in I$ and $\lambda \in [0, 1]$; If the inequality (2.1) is reversed, then the function $f(x)$ is said to be geometrically concave on I .

The notion of geometric convexity was introduced in [9]. For further development on convexity of functions, please refer to [10].

Lemma 2.1 ([10, Proposition 4.3]). *Let $f : I \subset (0, \infty) \rightarrow (0, \infty)$ be a twice differentiable function. The following assertions are equivalent:*

- (1) *The function f is geometrically convex;*
- (2) *The function $\frac{xf'(x)}{f(x)}$ is increasing;*
- (3) *The function f is geometrically convex if and only if*

$$x\{f(x)f''(x) - [f'(x)]^2\} + f(x)f'(x) \geq 0 \quad (2.2)$$

holds for all $x \in I$.

Lemma 2.2 ([13]). *Let $f : (a, b) \subseteq (0, \infty) \rightarrow (0, \infty)$ be a geometrically concave function. Then*

$$g(x) = \int_x^b f(t) dt \quad \text{and} \quad h(x) = \int_a^x f(t) dt$$

are also geometrically concave on (a, b) .

Lemma 2.3 ([12]). *Let $f : (a, b) \subseteq (0, \infty) \rightarrow (0, \infty)$ be monotonic and f^{-1} be the inverse of f . Then*

- (1) *if f is increasing and geometrically convex (or geometrically concave respectively), then f^{-1} is geometrically concave (or geometrically concave respectively);*

- (2) if f is decreasing and geometrically convex (or geometrically concave respectively), then f^{-1} is geometrically convex (or geometrically concave respectively).

3. PROOF OF THEOREM 1.1

We are now in a position to prove our main results.

Proof of Theorem 1.1. Let $f(t) = (1 - t^q)^{-1/p}$. Then an easy computation gives

$$\frac{tf'(t)}{f(t)} = \frac{q}{p} \cdot \frac{t^q}{1 - t^q} \quad (3.1)$$

and

$$\left(\frac{t^q}{1 - t^q} \right)' = \frac{qt^{q-1}}{(1 - t^q)^2}. \quad (3.2)$$

From (3.2) we see that the function $\frac{tf'(t)}{f(t)}$ is increasing. By Lemma 2.1, we deduce that the function $f(t)$ is geometrically convex. By Lemma 2.2, we find that the function $\arcsin_{p,q}(x)$ is geometrically convex.

On the other hand, let $h(x) = \arcsin_{p,q}(x)$ for $x \in (0, 1)$. Then

$$h'(x) = (1 - x^q)^{-1/p}$$

which is increasing. By Lemma 2.3, we derive that the function $\sin_{p,q}(x)$ is geometrically concave.

The rest may be proved similarly. \square

Proof of Corollary 1.1. This follows from Theorem 1.1 and Definition 2.1. \square

REFERENCES

- [1] G. D. Anderson, M. Vuorinen, and X.-H. Zhang, *Topics on special function III*, available online at <http://arxiv.org/abs/1209.1696>.
- [2] B. A. Bhayo and M. Vuorinen, *On generalized trigonometric functions with two parameters*, J. Approx. Theory **164** (2012), no. 10, 1415–1426; Available online at <http://dx.doi.org/10.1016/j.jat.2012.06.003>.
- [3] P. J. Bushell and D. E. Edmunds, *Remarks on generalized trigonometric functions*, Rocky Mountain J. Math. **42** (2012), no. 1, 25–57; Available online at <http://dx.doi.org/10.1216/RMJ-2012-42-1-25>.
- [4] D. E. Edmunds, P. Gurka, and J. Lang, *Properties of generalized trigonometric functions*, J. Approx. Theory **164** (2012), no. 1, 47–56; Available online at <http://dx.doi.org/10.1016/j.jat.2011.09.004>.
- [5] R. Klen, M. Vuorinen, and X.-H. Zhang, *Inequalities for the generalized trigonometric and hyperbolic functions*, available online at <http://arxiv.org/abs/1210.6749>.
- [6] J. Lang and D. E. Edmunds, *Eigenvalues, Embeddings and Generalized Trigonometric Functions*, Lecture Notes in Mathematics 2016, Springer-Verlag, 2011.
- [7] P. Lindqvist, *Some remarkable sine and cosine functions*, Ricerche Mat. **44** (1995), no. 2, 269–290 (1996).
- [8] P. Lindqvist and J. Peetre, *p-arclength of the q-circle*, Math. Student **72** (2003), no. 1-4, 139–145.
- [9] P. Montel, *Sur les fonctions convexes et les fonctions sousharmoniques*, J. Math. Pures Appl. **9** (1928), no. 7, 29–60.
- [10] C. P. Niculescu, *Convexity according to the geometric mean*, Math. Inequal. Appl. **3** (2000), no. 2, 155–167; Available online at <http://dx.doi.org/10.7153/mia-03-19>.
- [11] S. Takeuchi, *Generalized Jacobian elliptic functions and their application to bifurcation problems associated with p-Laplacian*, J. Math. Anal. Appl. **385** (2012), no. 1, 24–35; Available online at <http://dx.doi.org/10.1016/j.jmaa.2011.06.063>.

- [12] X.-M. Zhang and Y.-M. Chu, *New Discussion to Analytic Inequalities*, Harbin Institute of Technology Press, Harbin, China 2009. (Chinese).
- [13] X.-M. Zhang and Y.-M. Chu, *The geometrical convexity and concavity of integral for convex and concave functions*, Int. J. Mod. Math. **3** (2008), no. 3, 345–350.

(Jiang) DEPARTMENT OF INFORMATION ENGINEERING, WEIHAI VOCATIONAL UNIVERSITY, WEIHAI CITY, SHANDONG PROVINCE, 264210, CHINA
E-mail address: jackjwd@163.com

(Qi) DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300387, CHINA
E-mail address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com
URL: <http://qifeng618.wordpress.com>